

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Attorney Docket No.: 4318.224-US

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Bisgaard-Frantzen et al.

Confirmation No: 7527

Serial No.: 09/902,188

Group Art Unit: 1751

Filed: July 10, 2001

Examiner: To be assigned

For: Amylase Variants

PENDING CLAIMS

Commissioner for Patents
Washington, DC 20231FAX RECEIVED
DEC 19 2001
GROUP 1700

OFFICIAL

68. A variant of a parent alpha-amylase enzyme, wherein said parent alpha-amylase has an amino acid sequence which has at least 80% homology to SEQ ID NO:3, and wherein said variant comprises deletions at positions equivalent to positions 180 and 181 in SEQ ID NO:3 (using SEQ ID NO:3 for numbering).

69. The variant of claim 68, wherein said parent alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

70. The variant of claim 68, wherein said parent alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

71. The variant of claim 68, wherein said parent alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

72. The variant of claim 68, wherein said variant further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

73. An isolated alpha-amylase enzyme comprising an amino acid sequence having an amino acid sequence which has at least 80% homology to SEQ ID NO:3, modified by having deletions at positions equivalent to positions 180 and 181 in SEQ ID NO:3.

74. The alpha-amylase enzyme of claim 73, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to 349 and 428 in SEQ ID NO:3.

75. The alpha-amylase of claim 73, wherein said alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

76. The alpha-amylase of claim 73, wherein said alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

77. The alpha-amylase of claim 73, wherein said alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

78. A process for producing an alpha-amylase enzyme, said process comprising:

- a) cultivating a host cell having a nucleic acid sequence encoding an alpha-amylase enzyme, said alpha-amylase enzyme comprising an amino acid sequence having at least 80% homology to SEQ ID NO:3 and wherein said alpha-amylase enzyme is modified by having deletions at positions equivalent to positions 180 and 181 in SEQ ID NO:3, wherein said cultivating is performed under conditions conducive to produce the alpha-amylase enzyme, and
- b) recovering the alpha-amylase from the culture.

79. The process of claim 78, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

80. A DNA construct having a DNA sequence encoding an alpha-amylase enzyme comprising an amino acid sequence of SEQ ID NO:3 and further comprising deletions at positions equivalent to positions 180 and 181 in SEQ ID NO:3.

81. The DNA construct of claim 80, wherein said alpha-amylase enzyme further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

82. A recombinant expression vector carrying the DNA construct of claim 80.

83. A cell transformed with the DNA construct of claim 80.

84. A cell transformed with the recombination expression vector of claim 82.

85. A process of producing an alpha-amylase, said process comprising culturing the cell of claim 83 under conditions conducive for the production of the alpha-amylase and recovering the alpha-amylase from the culture.

86. A variant of a parent alpha-amylase enzyme, wherein said parent alpha-amylase has an amino acid sequence which has at least 80% homology to SEQ ID NO:3, and wherein said variant comprises deletions at positions equivalent to positions 179 and 181 in SEQ ID NO:3 (using SEQ ID NO:3 for numbering).

87. The variant of claim 86, wherein said parent alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

88. The variant of claim 86, wherein said parent alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

89. The variant of claim 86, wherein said parent alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

90. The variant of claim 86, wherein said variant further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

91. An isolated alpha-amylase enzyme comprising an amino acid sequence having an amino acid sequence which has at least 80% homology to SEQ ID NO:3, modified by having deletions at positions equivalent to positions 179 and 181 in SEQ ID NO:3.

92. The alpha-amylase enzyme of claim 91, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to 349 and 428 in SEQ ID NO:3.

93. The alpha-amylase of claim 91, wherein said alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

94. The alpha-amylase of claim 91, wherein said alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

95. The alpha-amylase of claim 91, wherein said alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

96. A process for producing an alpha-amylase enzyme, said process comprising:

- a) cultivating a host cell having a nucleic acid sequence encoding an alpha-amylase enzyme, said alpha-amylase enzyme comprising an amino acid sequence having at least 80% homology to SEQ ID NO:3 and wherein said alpha-amylase enzyme is modified by having deletions at positions equivalent to positions 179 and 181 in SEQ ID NO:3, wherein said cultivating is performed under conditions conducive to produce the alpha-amylase enzyme, and
- b) recovering the alpha-amylase from the culture.

97. The process of claim 96, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

98. A DNA construct having a DNA sequence encoding an alpha-amylase enzyme comprising an amino acid sequence of SEQ ID NO:3 and further comprising deletions at positions equivalent to positions 179 and 181 in SEQ ID NO:3.

99. The DNA construct of claim 98, wherein said alpha-amylase enzyme further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

100. A recombinant expression vector carrying the DNA construct of claim 98.

101. A cell transformed with the DNA construct of claim 98.

102. A cell transformed with the recombination expression vector of claim 100.

103. A process of producing an alpha-amylase, said process comprising culturing the cell of claim 101 under conditions conducive for the production of the alpha-amylase and recovering the alpha-amylase from the culture.

104. A variant of a parent alpha-amylase enzyme, wherein said parent alpha-amylase has an amino acid sequence which has at least 80% homology to SEQ ID NO:3, and wherein said variant comprises deletions at positions equivalent to positions 179 and 182 in SEQ ID NO:3 (using SEQ ID NO:3 for numbering).

105. The variant of claim 104, wherein said parent alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

106. The variant of claim 104, wherein said parent alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

107. The variant of claim 104, wherein said parent alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

108. The variant of claim 104, wherein said variant further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

109. An isolated alpha-amylase enzyme comprising an amino acid sequence having an amino acid sequence which has at least 80% homology to SEQ ID NO:3, modified by having deletions at positions equivalent to positions 179 and 182 in SEQ ID NO:3.

110. The alpha-amylase enzyme of claim 109, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to 349 and 428 in SEQ ID NO:3.

111. The alpha-amylase of claim 109, wherein said alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

112. The alpha-amylase of claim 109, wherein said alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

113. The alpha-amylase of claim 109, wherein said alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

114. A process for producing an alpha-amylase enzyme, said process comprising:

- a) cultivating a host cell having a nucleic acid sequence encoding an alpha-amylase enzyme, said alpha-amylase enzyme comprising an amino acid sequence having at least 80% homology to SEQ ID NO:3 and wherein said alpha-amylase enzyme is modified by having deletions at positions equivalent to positions 179 and 182 in SEQ ID NO:3, wherein said cultivating is performed under conditions conducive to produce the alpha-amylase enzyme, and
- b) recovering the alpha-amylase from the culture.

115. The process of claim 114, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

116. A DNA construct having a DNA sequence encoding an alpha-amylase enzyme comprising an amino acid sequence of SEQ ID NO:3 and further comprising deletions at positions equivalent to positions 179 and 182 in SEQ ID NO:3.

117. The DNA construct of claim 116, wherein said alpha-amylase enzyme further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

118. A recombinant expression vector carrying the DNA construct of claim 116.

119. A cell transformed with the DNA construct of claim 116.

120. A cell transformed with the recombination expression vector of claim 118.

121. A process of producing an alpha-amylase, said process comprising culturing the cell of claim 119 under conditions conducive for the production of the alpha-amylase and recovering the alpha-amylase from the culture.

122. A variant of a parent alpha-amylase enzyme, wherein said parent alpha-amylase has an amino acid sequence which has at least 80% homology to SEQ ID NO:3, and wherein said variant comprises deletions at positions equivalent to positions 180 and 182 in SEQ ID NO:3 (using SEQ ID NO:3 for numbering).

123. The variant of claim 122, wherein said parent alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

124. The variant of claim 122, wherein said parent alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

125. The variant of claim 122, wherein said parent alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

126. The variant of claim 122, wherein said variant further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

127. An isolated alpha-amylase enzyme comprising an amino acid sequence having an amino acid sequence which has at least 80% homology to SEQ ID NO:3, modified by having deletions at positions equivalent to positions 180 and 182 in SEQ ID NO:3.

128. The alpha-amylase enzyme of claim 127, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to 349 and 428 in SEQ ID NO:3.

129. The alpha-amylase of claim 127, wherein said alpha-amylase has an amino acid sequence which has at least 85% homology to SEQ ID NO:3.

130. The alpha-amylase of claim 127, wherein said alpha-amylase has an amino acid sequence which has at least 90% homology to SEQ ID NO:3.

131. The alpha-amylase of claim 127, wherein said alpha-amylase has an amino acid sequence which has at least 95% homology to SEQ ID NO:3.

132. A process for producing an alpha-amylase enzyme, said process comprising:

- a) cultivating a host cell having a nucleic acid sequence encoding an alpha-amylase enzyme, said alpha-amylase enzyme comprising an amino acid sequence having at least 80% homology to SEQ ID NO:3 and wherein said alpha-amylase enzyme is modified by having deletions at positions equivalent to positions 180 and 182 in SEQ ID NO:3, wherein said cultivating is performed under conditions conducive to produce the alpha-amylase enzyme, and
- b) recovering the alpha-amylase from the culture.

133. The process of claim 132, wherein said alpha-amylase enzyme is further modified by having amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

134. A DNA construct having a DNA sequence encoding an alpha-amylase enzyme comprising an amino acid sequence of SEQ ID NO:3 and further comprising deletions at positions equivalent to positions 180 and 182 in SEQ ID NO:3.

135. The DNA construct of claim 134, wherein said alpha-amylase enzyme further comprises amino acid substitutions of a cysteine at positions equivalent to positions 349 and 428 in SEQ ID NO:3.

136. A recombinant expression vector carrying the DNA construct of claim 134.

137. A cell transformed with the DNA construct of claim 134.

138. A cell transformed with the recombination expression vector of claim 136.

139. A process of producing an alpha-amylase, said process comprising culturing the cell of claim 137 under conditions conducive for the production of the alpha-amylase and recovering the alpha-amylase from the culture.